

Recombinant Human S100A6

Catalog#:P00170	Derived from E.coli
1 ····· Allower S100 Collins Disting Disting	

DESCRIPTION	Recombinant Human S100 Calcium Binding Protein A is produced by our E.coli expression system and the target gene encoding Met1-Gly90 is expressed with a 6His tag at the N-terminus. Accession#: P06703 Known as: S100A6; Protein S100-A6; Calcyclin; Growth factor-inducible protein 2A9; MLN 4; Prolactin receptor-associated protein; PRA;S100 calcium-binding protein A6; CACY	
FORMULATION	Supplied as a 0.2 µm filtered solution of 20mM Tris-HCl, 150mM NaCl, 20% Glycerol, 1mM EDTA, pH 8.0.	
SHIPPING	The product is shipped on dry ice/polar packs. Upon receipt, store it immediately at the temperature listed below.	
STORAGE	Store at \leq -70°C, stable for 6 months after receipt. Store at \leq -70°C, stable for 3 months under sterile conditions after opening. Please minimize freeze-thaw cycles.	
QUALITY CONTROLMol Mass:12.5kDaAP Mol Mass:12kDa, reducing conditions.Purity: Greater than 95% as determined by reducing SDS-PAGE. Endotoxin: Less than 0.1 ng/µg (1 EU/µg) as determined by LAL test.		
BACKGROUND	S100A6, also known as Protein S100-A6, Calcyclin, Growth factor-inducible protein 2A9, MLN 4, Prolactin receptor-associated protein, PRA, S100 calcium-binding protein A6 and CACY, is a member of the S100 family of proteins containing 2 EF-hand calcium-binding motifs. S100 proteins are a family of low molecular weight protein found in vertebrates and localized in the cytoplasm and/or nucleus of a wide range of cells. S100 proteins are	
ACKGROUND involved in a number of fundamental biological processes such as protein phosphorylation, transcription factors, the dynamics of cytoskeleton constituents, enzyme activities, cell growth and differentiation, the inflammatory response,cell cycle progression and differentiation, stimulation of Ca 2+-dependent insulin release, stimulation of prolactin secretion, and exocytosis. Chromosomal rearrangements and altered expression of this gene have been implicated in melanoma. S100A6 may function as calcium sensor and modulator, contributing to cellular calcium signaling. It may function by interacting with other proteins, such as TPR -containing proteins, and indirectly play a role in many physiological processes such as the reorganization of the actin cytoskeleton and in cell motility.		
kDa MK R 120 90 90 60 40 30 30 14 7 14		