


Recombinant Human EGFR

Catalog#:P01024 Derived from Human Cells

DESCRIPTION	<p>Recombinant Human Epidermal Growth Factor Receptor is produced by our Mammalian expression system and the target gene encoding Leu25-Ser645 is expressed with a Fc tag at the C-terminus.</p> <p>Accession#: P00533</p> <p>Known as: Epidermal growth factor receptor;Proto-oncogene c-ErbB-1;Receptor tyrosine-protein kinase erbB-1;EGFR;ERBB; ERBB1; HER1</p>
FORMULATION	Lyophilized from a 0.2 μm filtered solution of PBS, pH 7.4.
SHIPPING	<p>The product is shipped at ambient temperature.</p> <p>Upon receipt, store it immediately at the temperature listed below.</p>
STORAGE	<p>Lyophilized protein should be stored at <-20°C, though stable at room temperature for 3 weeks.</p> <p>Reconstituted protein solution can be stored at 4-7°C for 2-7 days.</p> <p>Aliquots of reconstituted samples are stable at < -20°C for 3 months.</p>
RECONSTITUTION	<p><i>Always centrifuge tubes before opening. Do not mix by vortex or pipetting.</i></p> <p><i>It is not recommended to reconstitute to a concentration less than 100μg/ml.</i></p> <p>Dissolve the lyophilized protein in distilled water.</p> <p>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.</p>
QUALITY CONTROL	<p>Mol Mass:95.7kDa AP Mol Mass:120-160kDa, reducing conditions.</p> <p>Purity: Greater than 95% as determined by reducing SDS-PAGE.</p> <p>Endotoxin: Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.</p>
BACKGROUND	<p>EGFR is a transmembrane glycoprotein that is a member of the protein kinase superfamily. This protein is a receptor for members of the epidermal growth factor family. EGFR is a cell surface protein that binds to epidermal growth factor. Binding of the protein to a ligand induces receptor dimerization and tyrosine autophosphorylation and leads to cell proliferation. Activates at least 4 major downstream signaling cascades including the RAS-RAF-MEK-ERK, PI3 kinase-AKT, PLCgamma-PKC and STATs modules. May also activate the NF-kappa-B signaling cascade. Also directly phosphorylates other proteins like RGS16, activating its GTPase activity and probably coupling the EGF receptor signaling to the G protein-coupled receptor signaling.</p>
 <p>SDS-PAGE</p> <p>kDa MK R</p> <p>120</p> <p>90</p> <p>60</p> <p>40</p> <p>30</p> <p>20</p> <p>14</p>	