

Tel: 400-968-6088 Fax: 010-56371281

Http://www.solarbio.com

## **Recombinant Human DKK-3**

Catalog#:P00769 Derived from Human Cells

DESCRIPTION	Recombinant Human Dickkopf- Related Protein 3 is produced by our Mammalian expression system and the target gene encoding Ala22- Ile350 is expressed with a 6His tag at the C-terminus.  Accession#: AAF02676.1  Known as: Dickkopf- Related Protein 3; Dickkopf-3; Dkk-3; hDkk-3; DKK3; REIC
FORMULATION	Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4.
SHIPPING	The product is shipped at ambient temperature. Upon receipt, store it immediately at the temperature listed below.
STORAGE	Lyophilized protein should be stored at<-20°C, though stable at room temperature for 3 weeks.  Reconstituted protein solution can be stored at 4-7°C for 2-7 days.  Aliquots of reconstituted samples are stable at < -20°C for 3 months.
RECONSTITUTION	Always centrifuge tubes before opening. Do not mix by vortex or pipetting. It is not recommended to reconstitute to a concentration less than 100µg/ml. Dissolve the lyophilized protein in distilled water. Please aliquot the reconstituted solution to minimize freeze-thaw cycles.
QUALITY CONTROL	Mol Mass:37.22kDa AP Mol Mass:73kDa, reducing conditions.  Purity: Greater than 95% as determined by reducing SDS-PAGE.  Endotoxin: Less than 0.1 ng/μg (1 EU/μg) as determined by LAL test.
BACKGROUND	Dickkopf-related protein 3 (DKK3) belongs to the DKK protein family including Dkk-1, 2, 3 and -4. DKK3 is a 350 amino acid secreted glycoprotein which is comprised of an N-terminal signal peptide and 2 conserved cysteine-rich domains that are separated by a 12 amino acid linker region. Dkk-3 also have one prokineticin domain. DKK3 is involved in embryonic development through its inhibition of the WNT signaling pathway. The Dkk family also includes Soggy, which is homologous to Dkk-3 but not to the other family members. Soggy has not been shown to inhibit Wnt signaling, and its role in the pathway is unclear.
kDa MK R	

