


## Recombinant Human IL-8 (C-6His)

Catalog#:P00132    Derived from Human Cells

<b>DESCRIPTION</b>	<p>Recombinant Human C-X-C Motif Chemokine 8/Interleukin 8 is produced by our Mammalian expression system and the target gene encoding Glu21-Ser99 is expressed with a 6His tag at the C-terminus.</p> <p>Accession#:P10145</p> <p>Known as: Interleukin-8; IL-8; C-X-C Motif Chemokine 8; CXCL8; Emotakin; Granulocyte Chemotactic Protein 1; GCP-1; Monocyte-Derived Neutrophil Chemotactic Factor; MDNCF; Monocyte-Derived Neutrophil-Activating Peptide; MONAP; NeutrophilActivating Protein 1; NAP-1</p>
<b>FORMULATION</b>	Lyophilized from a 0.2 µm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.
<b>SHIPPING</b>	<p>The product is shipped at ambient temperature.</p> <p>Upon receipt, store it immediately at the temperature listed below.</p>
<b>STORAGE</b>	<p>Lyophilized protein should be stored at ≤ -20°C, stable for one year after receipt.</p> <p>Reconstituted protein solution can be stored at 2-8°C for 2-7 days.</p> <p>Aliquots of reconstituted samples are stable at ≤ -20°C for 3 months.</p>
<b>RECONSTITUTION</b>	<p>Always centrifuge tubes before opening. Do not mix by vortex or pipetting.</p> <p>It is not recommended to reconstitute to a concentration less than 100µg/ml.</p> <p>Dissolve the lyophilized protein in distilled water.</p> <p>Please aliquot the reconstituted solution to minimize freeze-thaw cycles.</p>
<b>QUALITY CONTROL</b>	<p>Mol Mass: 10.1kDa    AP Mol Mass: 14kDa, reducing conditions.</p> <p>Purity: Greater than 95% as determined by reducing SDS-PAGE.</p> <p>Endotoxin: Less than 0.1 ng/µg (1 EU/µg) as determined by LAL test.</p>
<b>BACKGROUND</b>	<p>Interleukin-8 (IL-8) belongs to the neutrophil-specific CXC family of chemokines. It is one of the initial cytokines released from a variety of cell types, including T cells, endothelial cells and fibroblasts, in response to an inflammatory stimulus and acts by recruiting neutrophils, T-cells and basophils to the site of inflammation. Elevated Interleukin-8 levels are associated with the onset of a variety of disease states.</p>
<div style="display: flex; align-items: center;"> <div style="margin-right: 20px;"> <p><b>SDS-PAGE</b></p> </div>  </div>	