

Hexokinase

Cat: H8210

Specification: 500U

Storage: Store at -20°C, and it is stable for at least 2 years when stored properly. Solutions in water or citrate buffer have remained stable during repeated freezing and thawing for a period of 30 days.

Product Information

CAS: 9001-51-8


Enzyme Activity/Potency: 426U/mg

Synonym: ATP:D-hexose 6-phosphotransferase

Molecular Weight: 110 kDa (dimer) Hexokinase is a dimeric protein with two equal 55 kDa monomers.

Introduction

Glycolysis is the process occurring in almost all living organisms by which they metabolize D-glucose to generate energy and metabolic intermediates. In the first step of glycolysis, hexokinase phosphorylates the C6 position of D-glucose in the presence of ATP by the following reaction:



Yeast hexokinase has three isozymes, designated P-I, P-II, and glucokinase (Glk1). Each has distinctive properties. Yeast hexokinase P-II has both a catalytic and a regulatory function.

Several other hexoses can serve as substrates for hexokinase (relative reaction rates):

Glucose; 2-deoxy-2-fluoro-D-glucose; mannosamine; 5-thioglucose; 3-deoxy-3-aminoglucose; 1, 5-anhydro-D-glucitol, 1-thio-D-glucose.

Activators: Mg^{2+} (KM = 2.6 mM), catecholamine-related compounds.

Inhibitors: D-glucosamine ; D-mannose; D-xylose; 6-deoxy-D-glucose ; N-acetylmannosamine. Also sorbose-1-phosphate, polyphosphates, 6-deoxy-6-fluoroglucose, 2-C-hydroxy-methylglucose, lyxose, and thiol reactive compounds.

Hexokinase is used for the determination of D-glucose, D-fructose, and D-sorbitol in food or other biological materials. Hexokinase can also be used in the assay of glycosides that are convertible to glucose or fructose.

Unit Definition: One unit will phosphorylate 1.0 μmole of D-glucose per min at pH 7.6 at 25 °C.

Preparation Instructions:

Hexokinase is soluble in cold water (0.5-1.0 unit/ml) or citrate buffer, pH 7.0.

Precautions and Disclaimer:

For Laboratory Use Only. Not for drug, household or other uses.