

ssssssThioredoxin Reductase (TrxR) Activity Assay Kit

Note: Take two or three different samples for prediction before test.

Operation Equipment: Spectrophotometer

Cat No: BC1150

Size: 50T/48S

Product Composition: Before use, please carefully check whether the volume of the reagent is consistent with the volume in the bottle. If you have any questions, please contact Solarbio staff in time.

Reagent name	Size	Preservation Condition
Reagent I	Liquid 100 mL×1	2-8°C
Reagent II	Liquid 60 mL×1	2-8°C
Reagent III	Powder×2	-20°C
Reagent IV	Liquid 30 μL×1	-20°C

Solution Preparation:

1. Reagent III: The reagent was placed in a glass bottle, and before use, one bottle was dissolved in 3.33 mL distilled water, and the reagent was stored at -20°C for 2 weeks.
2. Reagent IV: Small volume, please centrifuge before use. Before clinical use, the reagent was diluted 10 times with anhydrous ethanol four times according to the number of samples.

Product Description:

Thioredoxin Reductase (TrxR) is a NADPH-dependent dimer selenase and includes FAD structure domain. TrxR belongs to pyridine nucleotide-disulfide REDOX enzyme, and form thioredoxin system with thioredoxin and NADPH. The activity of TrxR is similar with GR. TrxR could catalyzes GSSG reduct to GSH, which is the key enzyme in glutathione REDOX cycle.

TrxR catalyzes NADPH to reductDTNB form TNB and NADP⁺, TNB has a absorbance at 412 nm, but reduced glutathione reacts with DTNB to form TNB, so the 2-Vinylpyridine in this kit can inhibit reduced glutathione in sample. The activity of TrxR can be calculated by detecting increase rate of TNB at 412 nm.

Reagents and Equipment Required but Not Provided:

Spectrophotometer, low temperature centrifuge, adjustable pipette, 1 mL glass cuvette and distilled water.

Procedure

I. Sample preparation:

1. Tissue:

Add 1 mL of Reagent I into 0.1 g of plant or animal tissue, fully grinding on ice. Centrifuge at 10000 rpm for 10 minutes at 4°C, take the supernatant and place it on ice for test. Before test, mix supernatant and Reagent IV at a ratio of 50:1(add 2 μL of Reagent IV to 100 μL of supernatant), water bath at 37°C for 30

minutes, then keep on ice for test.

2. Bacteria/cell:

Suggested 5 million with 1 mL of Reagent I, Splitting bacteria and cell with ultrasonic (ice bath, power 300W, work time 3 s, interval 7 s, for 3 min). Centrifuge at 10000 rpm for 10 minutes at 4°C, take the supernatant on ice for test. Before test, mix supernatant and Reagent IV at a ratio of 50:1 (add 2 μL of Reagent IV to 100 μL of supernatant), water bath at 37°C for 30 minutes, then keep on ice for test.

II. Procedure:

1. Preheat spectrophotometer for 30 minutes, adjust the wavelength to 412 nm, set the zero with distilled water.
2. Preheat Reagent I at 37°C (mammal), 25°C (other) in water bath for 30 minutes.
3. Blank tube: take a 1 mL glass cuvette, add 100 μL of Reagent II, 100 μL of Reagent III, 800 μL of Reagent I, mix them quickly, and then measure the absorbance at 412 nm for 10s. Take out the absorbance at 412 nm in a 37°C water bath for 5 min and record it as A1 and A2. Calculate $\Delta A_B = A_2 - A_1$.
4. Measuring tube: take a 1 mL glass cuvette, add 100 μL of Reagent II, 100 μL of Reagent III, 700 μL of Reagent I, 100 μL of supernatant, mix it quickly, and then measure the absorbance at 412 nm for 10s, take out the absorbance at 412 nm quickly in a 37°C water bath for 5 min, and record it as A3 and A4. $\Delta A_T = A_4 - A_3$.

III. Calculation:

1. Protein concentration:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the production of 1 μmol of TNB at 37°C (mammal), 25°C (other) per minute every mg of protein.

$$\text{TrxR (U/mg prot)} = [\Delta A(T) - \Delta A(B)] \div (\epsilon \times d) \times V_{rv} \div (V_s \times C_{pr}) \div T = 0.147 \times [\Delta A(T) - \Delta A(B)] \div C_{pr}$$

2. Sample weight:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the production of 1 μmol of TNB at 37°C (mammal), 25°C (other) per minute every gram of sample.

$$\text{TrxR (U/g)} = [\Delta A(T) - \Delta A(B)] \div (\epsilon \times d) \times V_{rv} \div (V_s \div V_{sv} \times W) \div T = 0.147 \times [\Delta A(T) - \Delta A(B)] \div W$$

3. Cells or bacteria:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the production of 1 μmol of TNB at 37°C (mammal), 25°C (other) per minute every 10⁴ cell.

$$\text{TrxR (U/10}^4 \text{ cell)} = [\Delta A(T) - \Delta A(B)] \div (\epsilon \times d) \times V_{rv} \div (N \div V_{sv} \times V_s) \div T = 0.147 \times [\Delta A(T) - \Delta A(B)] \div N$$

ϵ : TNB molar extinction coefficient, 1.36×10^4 L/mol/cm;

d : Light path of cuvette, 1 cm;

V_{rv} : Total reaction volume, 1000 μL = 0.001 L;

V_s : Supernatant volume (mL), 0.1 mL;

C_{pr} : Sample protein concentration (mg/mL); need to detect separately, suggest use PC0020, BCA Protein Assay Kit;

T: Reaction time (min), 5 minutes;

W: Sample weight(g);

Vsv: Extraction volume, 1 mL;

N: Amount of cells,10⁴.

Note:

1. Dilute 5 times with distilled water when detecting mammalian tissue and blood samples, detect quickly as soon as possible.
2. Because the extract solution contains a certain concentration of protein (about 0.1 mg/mL), the protein content of the extract solution itself needs to be subtracted when determining the protein concentration of the sample.

Experimental instances:

1. Take 0.1g of Chinese rose petals, add 1mL of extract solution, fully grinding on ice. Centrifuge at 10000 rpm for 10 minutes at 4°C, take the supernatant and place it on ice for test according to the measured steps. Calculate $\Delta A_T = A_4 - A_3 = 0.763 - 0.716 = 0.047$, $\Delta A_B = A_2 - A_1 = 0.082 - 0.064 = 0.018$, calculate the enzyme activity according to sample weight:

$$\text{TrxR (U/g weight)} = 147 \times (\Delta A_T - \Delta A_B) \div W = 4.26 \text{ U/g weight.}$$

2. Take 0.1g of liver, add 1mL of extract solution, fully grinding on ice. Centrifuge at 10000 rpm for 10 minutes at 4°C, take the supernatant and place it on ice for test according to the measured steps. Calculate $\Delta A_T = A_4 - A_3 = 1.508 - 0.4 = 1.108$, $\Delta A_B = A_2 - A_1 = 0.082 - 0.064 = 0.018$, calculate the enzyme activity according to sample weight:

$$\text{TrxR (U/g weight)} = 147 \times (\Delta A_T - \Delta A_B) \div W \times 2 \text{ (dilution ratio)} = 3204.6 \text{ U/g weight.}$$

Recent Product citations

[1] Li B, Li D, Jing W, et al. Biogenic selenium and its hepatoprotective activity[J]. Scientific reports, 2017, 7(1): 1-11.

[2] Zhang L, Fan J, He J, et al. Regulation of ROS–NF-κB axis by tuna backbone derived peptide ameliorates inflammation in necrotizing enterocolitis[J]. Journal of cellular physiology, 2019, 234(8): 14330-14338.

Related products:

BC1170/ BC1175	Reduced Glutathione (GSH) Assay Kit
BC1180/ BC1185	Oxidized Glutathione (GSSG) Assay Kit
BC1190/ BC1195	Glutathione Peroxidase (GPX) Assay Kit
BC0350/ BC0355	Glutathione S-transferase(GST) Activity Assay Kit