

## Fructose-1, 6-diphosphate (FDP) Activity Assay Kit

**Note:** The reagents have been changed, so please be aware of and follow this instruction strictly.

**Operation Equipment:** Spectrophotometer

**Cat No:** BC2240

**Size:** 50T/24S

### Components:

**Extract solution I:** Liquid 30 mL×1. Store at 2-8°C.

**Extract solution II:** Liquid 5 mL×1. Store at 2-8°C.

**Reagent I:** Liquid 20 mL×1. Store at 2-8°C.

**Reagent II:** Liquid×2. Store at 2-8°C. Dissolve with 0.5 mL of distilled water before use.

Unused reagents can be stored at 20°C for up to 4 weeks, avoiding repeated freezing and thawing..

**Reagent III:** Liquid 15 mL×1. Store at 2-8°C.

**Reagent IV:** Liquid 40 mL×1. Store at 2-8°C.

**Standard:** Powder×1. Store at 2-8°C. Dissolve with 1.176 mL of distilled water before use to form 50 μmol/mL FDP standard solution. The standard can be stored at 2-8°C for 4 weeks.

### Product Description:

Fructose-1,6-diphosphate (FDP) is an important intermediate product in the glycolysis process. It can regulate a variety of enzymes, improve cell energy metabolism, increase energy utilization, anti-arrhythmia and anti-tissue peroxidation. FDP is widely used in clinical medicine.

Aldolase catalyzes the cleavage of fructose 1,6-diphosphate. The product reacts with 2,4-dinitrophenylhydrazine in acid medium to form 2,4-dinitrophenylhydrazone, which is dark red in alkaline solution and has a characteristic absorption peak at 540 nm.

### Reagents and Equipment Required but Not Provided:

spectrophotometer, desk centrifuge, adjustable transferpettor, water bath /incubator, 1 mL glass cuvette, mortar / homogenizer, Ultrasonic crusher, ice and distilled water.

### Procedure:

#### I. Sample preparation:

##### 1) Tissue

According to the tissue weight (g): the volume of the extract (mL) is 1:5 ~ 10. Suggest adding 1 mL of Extract solution I to 0.1 g of tissue, fully homogenize on ice bath. Centrifuge at 12000 ×g for 10 minutes at 4°C. Add 0.15mL Extract solution II slowly to 0.8 mL supernatant. **Blend slowly until no bubbles**, centrifuge at 12000 ×g for 10 minutes at 4°C. Then take supernatant for test.

##### 2) Bacteria or cells

According to the Bacteria or cells ( $10^4$ ): the volume of the extract (mL) is 500~1000:1. Suggest add 1mL

of Extract solution to 5 million of bacteria or cells. Use ultrasonication to splitting bacteria and cells

(placed on ice, ultrasonic power 300W, working time 3 seconds, interval 7 seconds, total time for 3 min). Centrifuge at 12000 ×g for 10 minutes at 4°C. Add 0.15mL Extract solution II slowly to 0.8 mL supernatant. **Blend slowly until no bubbles**, centrifuge at 12000 ×g for 10 minutes at 4°C. Then take supernatant for test.

### 3) Liquid:

Add 1mL Extract solution I to 100μL liquid sample, centrifuge at 12000 ×g for 10 minutes at 4°C. Add 0.15mL Extract solution II slowly to 0.8 mL supernatant. **Blend slowly until no bubbles**, centrifuge at 12000 ×g for 10 minutes at 4°C. Then take supernatant for test.

### Determination procedure:

1. Preheat spectrophotometer for 30 minutes, adjust the wavelength to 540 nm, set zero with distilled water.
2. 50 μmol/ml fructose-1,6-diphosphate standard solution is diluted to 1.5625, 0.78125, 0.39, 0.2 and 0.1 μmol/ml standard solution with distilled water.
3. Sampling table:

Reagent name (μL)	Control tube (A <sub>C</sub> )	Test tube (A <sub>T</sub> )	Blank tube (A <sub>B</sub> )	Standard tube (A <sub>S</sub> )
Sample	100	100	-	-
Distilled water	-	-	100	-
Standard solution			-	100
Reagent I	220	200	220	200
Reagent II	-	20	-	20
Mix well, react accurately at 37°C for 2 h				
Reagent III	200	200	200	200
Mix well, react accurately at 37°C for 20 min				
Reagent IV	500	500	500	500
Mix well, react accurately at 37°C for 10 min				
The absorbance value at 540 nm is measured in 1 ml glass cuvette and recorded as A <sub>C</sub> , A <sub>T</sub> , A <sub>B</sub> , A <sub>S</sub> , respectively. Calculate $\Delta A = A_T - A_C$ , $\Delta A_S = A_S - A_B$ . The blank tube only needs to be tested 1-2 times.				

### II. Calculation:

1. According to concentration of standard solution and  $\Delta A_S$  to create the standard curve, take standard solution as X-axis,  $\Delta A_S$  as Y-axis. Take  $\Delta A$  into the equation to obtain x (μmol/ml).

#### 2. Calculation:

(1) sample weight

$$\text{FDP } (\mu\text{g/g weight}) = x \times (V_{\text{su}} + V_{\text{exII}}) \times M \div (W \times V_{\text{su}} \div V_{\text{exI}}) = 403.75X \div W$$

(2) The number of bacteria or cells

$$\text{FDP } (\mu\text{g}/10^4 \text{ cell}) = x \times (V_{\text{su}} + V_{\text{exII}}) \times M \div (N \times V_{\text{su}} \div V_{\text{exI}}) = 403.75X \div N$$

(3) Liquid:

$$\text{FDP } (\mu\text{g/mL}) = x \times (V_{\text{su}} + V_{\text{exII}}) \times M \div (V_{\text{L}} \times V_{\text{su}} \div (V_{\text{exI}} + V_{\text{L}})) = 4441.25x$$

$V_{\text{su}}$ : Supernatant volume of extraction, 0.8mL

$V_{\text{exII}}$ : Extract solution II volume, 0.16mL

M: Molecular weight of fructose-1,6-diphosphate, 340

$V_{\text{exI}}$ : Extract solution I volume, 1mL

W: sample weight, g

N: Cell amount, 10 thousand cells as unit

$V_{\text{L}}$ : liquid sample volume, 0.1mL.

**Note:**

1. If  $\Delta A > 0.5$ , please dilute the sample with water to appropriate concentration, multiply dilute times in the formula.

**Related Products:**

BC2270/BC2275 Fructose-bisphosphate aldolase(FBA) Activity Assay Kit

BC2250/BC2255 Phosphoglycerate Kinase(PGK) Activity Assay Kit