

Cellulase (CL) Activity Assay Kit

Note: Take two or three different samples for prediction before test.

Operation Equipment: Spectrophotometer

Catalog Number: BC2540

Size: 50T/24S

Components:

Extract reagent: 30mL×1, storage at 2-8°C.

Reagent I: 4mL×1, storage at 2-8°C.

Reagent II: 10mL×1, storage at 2-8°C.

Reagent III: 13mL×1, storage at 2-8°C.

Standard: powder ×1, storage at 2-8°C. Containing 10 mg of anhydrous glucose (loss on drying < 0.2%). Add 1 mL of distilled water to dissolve it for standby before use. It can be stored for two week at 4°C, or it can be stored for a longer time with saturated benzoic acid solution.

Product Description:

Cellulase (EC 3.2.1.4) exists in bacteria, fungi and animals, which can catalyze cellulose degradation. It is a type of enzyme preparation that can be widely used in the fields of medicine, food, cotton spinning, environmental protection and renewable resource utilization.

The 3,5-dinitrosalicylic acid method is used to determine the reducing sugar content of cellulose catalyzed by CL.

Reagents and Equipment Required but Not Provided:

Spectrophotometer, water bath, adjustable transferpeltor, mortar/homogenizer, centrifuge, 1mL glass cuvette, ice and distilled water.

Sample preparation:

1. Bacteria or cells: Collect the bacteria or cells into a centrifuge tube, discard the supernatant after centrifugation; add 1 mL of Extract reagent for every 5 million bacteria or cells, and break the bacteria or cells with an ultrasonic ice bath (power 200w, ultrasonic 3 seconds, interval 10 seconds, repeat 30 times); Centrifugate at 8000g and 4 °C for 10min, take the supernatant and place on ice for testing.
2. Plant and animal tissues: Weigh about 0.1 g of sample, add 1 mL of Extract reagent and fully grind. Centrifugate at 8000g and 4°C for 10 min, the supernatants as samples to be tested.

Procedure:

1. Preheat spectrophotometer for 30min, adjust the wavelength to 540 nm and set the counter to zero with distilled water.
2. Prepared standard: The standard was diluted with distilled water to 1, 0.8, 0.6, 0.4, 0.2, 0.1, 0 mg/mL..
3. Add reagent to a 1.5 mL EP tube:

Reagent name (μL)	Control tube (Ac)	Test tube (At)	Standard tube (As)
Reagent I	50	50	-
Reagent II	200	200	-
Distilled water	50	50	-
Sample		50	-
Boiled sample	50		-
Mix well, and react accurately in water bath at 40°C for 30min. after taking out, put it in boiling water and boil for 15min immediately to get the saccharification solution.			
Saccharification solution	50	50	-
Standard solution	-	-	50
Reagent III	150	150	150
Mix well, boil for 15min in a boiling water bath and cool.			
Distilled water	1050	1050	1050

Mix well, set the counter to zero with distilled water, and measure the absorbance at 540 nm, and calculate $\Delta A = A_t - A_c$. $\Delta A_s = A_s - A(0 \text{ mg/mL})$. Standard curves only need to be done 1-2 times.

Calculation:

1. Standard curve

A standard curve was created from the absorbance (x, ΔA_s) and concentration (y, mg/mL) of the standard tube, and ΔA_t (x, ΔA_t) was brought into the standard curve to calculate the amount of product y (mg/mL) generated by the sample.

2. Calculation

(1) Protein concentration:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme that catalyzes the production of 1μg glucose per minute in the reaction system every milligram tissue protein

$$CL (U/mg \text{ prot}) = 1000 \times y \times V_{rv} \div (V_s \times C_{pr}) \div T = 233y \div C_{pr}$$

(2) Sample weight:

Unit definition: One unit of enzyme activity is defined as that one gram tissue catalyzes the production of 1μg glucose per min in the reaction system.

$$CL (U/g) = 1000 \times y \times V_{rv} \div (V_s \times W \div V_e) \div T = 233y \div W$$

(3) Cells or bacteria:

Unit definition: One unit of enzyme activity is defined as that 10^4 cells or bacteria catalyzes the production of 1μg glucose in the reaction system per min.

$$CL (U/10^4 \text{ cell}) = 1000 \times y \times V_{rv} \div (500 \times V_s \div V_e) \div T = 0.467 \times y$$

1000: 1mg/mL = 1000μg/mL

V_{rv} : Total volume of reaction system, 0.35mL.

V_s : sample volume added, 0.05mL;

V_e : volume used in the extraction solution, 1mL;

Cpr: sample protein concentration, mg/mL;

W: Fresh weight of sample, g;

T: React time, 30min.

500: the number of cells or bacteria, 5 million

Recent Product Citations:

Guo Q, Du G, Qi H, et al. A nematicidal tannin from *Punica granatum* L. rind and its physiological effect on pine wood nematode (*Bursaphelenchus xylophilus*)[J]. *Pesticide biochemistry and physiology*, 2017, 135: 64-68.

References:

Faria M L, Kolling D, Camassola M, et al. Comparison of *Penicillium echinulatum* and *Trichoderma reesei* cellulases in relation to their activity against various cellulosic substrates[J]. *Biores. Technol*, 2008, 99: 1417-1424.

Related Products:

BC0340/BC0345 Glucogen Content Assay Kit

BC2450/BC2455 Plant Tissue Fructose Content Assay Kit

BC2510/BC2515 Trehalase Activity Assay Kit